

REPRESENTATIONS OF $SL_2(\mathbb{R})$ AND NEARLY HOLOMORPHIC MODULAR FORMS

AMEYA PITALE, ABHISHEK SAHA, AND RALF SCHMIDT

ABSTRACT. In this semi-expository note, we give a new proof of a structure theorem due to Shimura for nearly holomorphic modular forms on the complex upper half plane. Roughly speaking, the theorem says that the space of all nearly holomorphic modular forms is the direct sum of the subspaces obtained by applying appropriate weight-raising operators on the spaces of holomorphic modular forms and on the one-dimensional space spanned by the weight 2 nearly holomorphic Eisenstein series.

While Shimura's proof was classical, ours is representation-theoretic. We deduce the structure theorem from a decomposition for the space of *n-finite* automorphic forms on $SL_2(\mathbb{R})$. To prove this decomposition, we use the mechanism of *category* \mathcal{O} and a careful analysis of the various possible indecomposable submodules. It is possible to achieve the same end by more direct methods, but we prefer this approach as it generalizes to other groups.

This note may be viewed as the toy case of our paper [6], where we prove an analogous structure theorem for vector-valued nearly holomorphic Siegel modular forms of degree two.

1. NEARLY HOLOMORPHIC FUNCTIONS

Let \mathbb{H}_1 be the complex upper half plane. Let $N^p(\mathbb{H}_1)$ be the space of functions $f : \mathbb{H}_1 \rightarrow \mathbb{C}$ of the form

$$f(\tau) = \sum_{j=0}^p f_j(\tau)y^{-j}, \quad \tau = x + iy,$$

where f_0, \dots, f_p are holomorphic functions on \mathbb{H}_1 . Any element of the space $N(\mathbb{H}_1) = \cup_{p=0}^{\infty} N^p(\mathbb{H}_1)$ is called a nearly holomorphic function on \mathbb{H}_1 . It is an exercise to show that

$$\sum_{j=0}^p f_j(\tau)y^{-j} = 0 \quad \iff \quad f_j = 0 \text{ for all } j = 0, \dots, p. \quad (1)$$

Hence, the holomorphic coefficients of a nearly holomorphic function are uniquely determined.

If f is a nearly holomorphic function, and if there exists a non-zero real number r such that $f(\tau+r) = f(\tau)$ for all $\tau \in \mathbb{H}_1$, then the holomorphic coefficients f_j of f exhibit the same translation invariance; this follows from (1). Each f_j therefore admits a Fourier expansion $f_j(\tau) = \sum a_j(n)e^{2\pi in\tau/r}$. It follows that f admits a Fourier expansion whose coefficients are polynomials in y^{-1} .

A.S. is partially supported by EPSRC grant EP/L025515/1. A.P. and R.S. are supported by NSF grant DMS-1100541.

For an integer k , we define the weight k slash operator on functions $f : \mathbb{H}_1 \rightarrow \mathbb{C}$ in the usual way:

$$(f|_k g)(\tau) = (c\tau + d)^{-k} f\left(\frac{a\tau + b}{c\tau + d}\right), \quad g = \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in \mathrm{SL}_2(\mathbb{R}).$$

Let Γ be a congruence subgroup of $\mathrm{SL}_2(\mathbb{Q})$. Let $N_k^p(\Gamma)$ denote the space of functions $F : \mathbb{H}_1 \rightarrow \mathbb{C}$ such that

- (1) $F \in N^p(\mathbb{H}_1)$;
- (2) $F|_k \gamma = F$ for all $\gamma \in \Gamma$;
- (3) F satisfies the cusp condition. (This notion is defined in terms of Fourier expansions just as in the case of holomorphic modular forms; see, e.g., §2.1 of [5].)

We denote by $N_k^p(\Gamma)^\circ$ the subspace of functions that vanish at every cusp. The space $N_k(\Gamma) = \cup_{p=0}^\infty N_k^p(\Gamma)$ is the space of *nearly holomorphic modular forms* with respect to Γ , and $N_k(\Gamma)^\circ = \cup_{p=0}^\infty N_k^p(\Gamma)^\circ$ is the space of *nearly holomorphic cusp forms*. Evidently, $M_k(\Gamma) := N_k^0(\Gamma)$ is the usual space of holomorphic modular forms of weight k with respect to Γ , and $S_k(\Gamma) := N_k^0(\Gamma)^\circ$ is the subspace of cusp forms. Nearly holomorphic modular forms occur naturally as special values of Eisenstein series and thus their arithmetic properties imply arithmetic properties for various L -functions via the theory of Rankin-Selberg type integrals. We refer the reader to the introduction of [6] for further remarks in this direction.

For an integer k , we define the classical Maass weight raising and lowering operators R_k, L_k on the space of smooth functions on \mathbb{H}_1 by

$$R_k = \frac{k}{y} + 2i \frac{\partial}{\partial \tau}, \quad L_k = -2iy^2 \frac{\partial}{\partial \bar{\tau}}, \quad (2)$$

where $\frac{\partial}{\partial \tau} = \frac{1}{2}(\frac{\partial}{\partial x} - i \frac{\partial}{\partial y})$ and $\frac{\partial}{\partial \bar{\tau}} = \frac{1}{2}(\frac{\partial}{\partial x} + i \frac{\partial}{\partial y})$ are the usual Wirtinger derivatives. Also define an operator Ω_k by

$$\Omega_k = \frac{1}{4}k^2 + \frac{1}{2}R_{k-2}L_k + \frac{1}{2}L_{k+2}R_k. \quad (3)$$

A calculation shows that

$$\Omega_k = y^2 \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) - 2iky \frac{\partial}{\partial \bar{\tau}} + \frac{k}{2} \left(\frac{k}{2} - 1 \right). \quad (4)$$

The following lemma is readily verified.

Lemma 1.1. *Let k be an integer, and p be a non-negative integer. Let Γ be a congruence subgroup of $\mathrm{SL}_2(\mathbb{Q})$.*

- (1) R_k induces maps $N_k^p(\Gamma) \rightarrow N_{k+2}^{p+1}(\Gamma)$ and $N_k^p(\Gamma)^\circ \rightarrow N_{k+2}^{p+1}(\Gamma)^\circ$.
- (2) L_k induces maps $N_k^p(\Gamma) \rightarrow N_{k-2}^{p-1}(\Gamma)$ and $N_k^p(\Gamma)^\circ \rightarrow N_{k-2}^{p-1}(\Gamma)^\circ$.
- (3) Ω_k induces endomorphisms of $N_k^p(\Gamma)$ and of $N_k^p(\Gamma)^\circ$.

Here, we understand $N_k^p(\Gamma) = N_k^p(\Gamma)^\circ = 0$ for $p < 0$.

Henceforth, we drop the subscripts and let R, L , and Ω denote the operators on $\bigoplus_k N_k(\Gamma)$ whose restrictions to $N_k(\Gamma)$ are given by R_k, L_k , and Ω_k , respectively.

Lemma 1.2. *For any integer k and non-negative integer p , the space $N_k^p(\Gamma)$ is finite-dimensional.*

Proof. This is well known for $p = 0$, since $N_k^0(\Gamma) = M_k(\Gamma)$ is simply the space of holomorphic modular forms of weight k . For $p > 0$ there is an exact sequence

$$0 \longrightarrow M_k(\Gamma) \longrightarrow N_k^p(\Gamma) \xrightarrow{L} N_{k-2}^{p-1}(\Gamma).$$

Hence the assertion follows by induction on p . \square

The following well-known fact will be important for our arguments further below. (For a proof, see Theorem 2.5.2 of [5].)

Lemma 1.3. $S_k(\Gamma) = 0$ if $k \leq 0$, and $M_k(\Gamma) = 0$ if $k < 0$. The space $M_0(\Gamma)$ consists of the constant functions.

2. REPRESENTATIONS OF $\mathrm{SL}_2(\mathbb{R})$ AND DIFFERENTIAL OPERATORS

To reinterpret elements of $N_k^p(\Gamma)$ as functions on $\mathrm{SL}_2(\mathbb{R})$, we recall the basic representation theory of this group. Let $\mathfrak{g} = \mathfrak{sl}_2(\mathbb{R})$ be the Lie algebra of $\mathrm{SL}_2(\mathbb{R})$, consisting of all 2×2 real matrices with trace zero. Let $\mathfrak{g}_{\mathbb{C}} = \mathfrak{sl}_2(\mathbb{C})$ be its complexification. The elements

$$H = -i \begin{bmatrix} 0 & 1 \\ -1 & 0 \end{bmatrix}, \quad R = \frac{1}{2} \begin{bmatrix} 1 & i \\ i & -1 \end{bmatrix}, \quad L = \frac{1}{2} \begin{bmatrix} 1 & -i \\ -i & -1 \end{bmatrix} \quad (5)$$

of $\mathfrak{g}_{\mathbb{C}}$ satisfy the relations $[H, R] = 2R$, $[H, L] = -2L$ and $[R, L] = H$. The *Casimir element* is the element in the universal enveloping algebra $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ given by

$$\Omega = \frac{1}{4}H^2 + \frac{1}{2}RL + \frac{1}{2}LR. \quad (6)$$

Then Ω lies in the center \mathcal{Z} of $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$, and it is known that $\mathcal{Z} = \mathbb{C}[\Omega]$.

Let $K = \mathrm{SO}(2)$ be the standard maximal compact subgroup of $\mathrm{SL}_2(\mathbb{R})$, consisting of all elements $r(\theta) = \begin{bmatrix} \cos(\theta) & \sin(\theta) \\ -\sin(\theta) & \cos(\theta) \end{bmatrix}$ with $\theta \in \mathbb{R}$. By ‘‘representation of $\mathrm{SL}_2(\mathbb{R})$ ’’ we mean a (\mathfrak{g}, K) -module. In such a module (π, V) , we say a non-zero $v \in V$ has weight k if

$$\pi(r(\theta))v = e^{ik\theta}v \quad \text{for } \theta \in \mathbb{R},$$

or equivalently, $\pi(H)v = kv$. In an irreducible representation, all weights have the same parity, and every weight occurs at most once. The operator $\pi(R)$ raises the weight by 2, and the operator $\pi(L)$ lowers the weight by 2. The *weight structure* of an irreducible representation is the list of weights, written in order. The following is the complete list of irreducible, admissible (\mathfrak{g}, K) -modules.

- (1) *Finite-dimensional representations.* For a positive integer p , let \mathcal{F}_p be the irreducible finite-dimensional representation of $\mathrm{SL}_2(\mathbb{R})$ with weight structure $[-p+1, -p+3, \dots, p-3, p-1]$. Hence $\dim \mathcal{F}_p = p$.
- (2) *Discrete series representations.* For a positive integer p we denote by $\mathcal{D}_{p,+}$ the discrete series representation of $\mathrm{SL}_2(\mathbb{R})$ with weight structure $[p+1, p+3, \dots]$. Similarly, let $\mathcal{D}_{p,-}$ be the discrete series representation of $\mathrm{SL}_2(\mathbb{R})$ with weight structure $[\dots, -p-3, -p-1]$. Hence, p is not the minimal weight of $\mathcal{D}_{p,+}$, but the Harish-Chandra parameter.
- (3) *Limits of discrete series.* Let $\mathcal{D}_{0,+}$ be the irreducible representation of $\mathrm{SL}_2(\mathbb{R})$ with weight structure $[1, 3, 5, \dots]$, and let $\mathcal{D}_{0,-}$ be the irreducible representation of $\mathrm{SL}_2(\mathbb{R})$ with weight structure $[\dots, -5, -3, -1]$. Formally these representations look like members of the discrete series, but they are not square-integrable.
- (4) *Principal series representations.* Their weight structure is either $2\mathbb{Z}$ or $2\mathbb{Z} + 1$. For our purposes, all we need to know about principal series representations is that the operators R and L act injectively on such a (\mathfrak{g}, K) -module.

Functions on $\mathrm{SL}_2(\mathbb{R})$ and functions on \mathbb{H}_1 . Let $W(k)$ be the space of smooth functions $\Phi : \mathrm{SL}_2(\mathbb{R}) \rightarrow \mathbb{C}$ with the property $\Phi(gr(\theta)) = e^{ik\theta}\Phi(g)$ for all $\theta \in \mathbb{R}$ and $g \in \mathrm{SL}_2(\mathbb{R})$. These are the vectors of weight k under the right translation action on the space of smooth functions. The operator R induces a map $W(k) \rightarrow W(k+2)$, and L induces a map $W(k) \rightarrow W(k-2)$. Let W be the space of smooth functions on \mathbb{H}_1 . For $\Phi \in W(k)$ we define an element $\tilde{\Phi} \in W$ by

$$\tilde{\Phi}(x+iy) = y^{-k/2} \Phi\left(\begin{bmatrix} 1 & x \\ & 1 \end{bmatrix} \begin{bmatrix} y^{1/2} & \\ & y^{-1/2} \end{bmatrix}\right). \quad (7)$$

It is straightforward to verify that

$$(\tilde{\Phi}|_k g)(i) = \Phi(g) \quad \text{for all } g \in \mathrm{SL}_2(\mathbb{R}). \quad (8)$$

The map $\Phi \mapsto \tilde{\Phi}$ establishes an isomorphism $W(k) \cong W$.

Lemma 2.1. *Let R, L, Ω be the operators on W defined in (2) and (4). Then the diagrams*

$$\begin{array}{ccccc} W(k) & \xrightarrow{\sim} & W & & W(k) & \xrightarrow{\sim} & W & & W(k) & \xrightarrow{\sim} & W \\ L \downarrow & & \downarrow L & & R \downarrow & & \downarrow R & & \Omega \downarrow & & \downarrow \Omega \\ W(k-2) & \xrightarrow{\sim} & W & & W(k+2) & \xrightarrow{\sim} & W & & W(k) & \xrightarrow{\sim} & W \end{array}$$

are commutative.

Proof. The assertions for R and L follow from straightforward calculations. The assertion for Ω then follows from (3) and (6). \square

The previous lemma is about smooth functions only and does not involve any transformation properties. If $\Phi \in W(k)$ satisfies $\Phi(\gamma g) = \Phi(g)$ for all $g \in \mathrm{SL}_2(\mathbb{R})$ and all elements γ of a congruence subgroup Γ , then it follows from (8) that $\tilde{\Phi}|_k \gamma = \tilde{\Phi}$ for all $\gamma \in \Gamma$. Conversely, given a smooth function f on \mathbb{H}_1 satisfying $f|_k \gamma = f$ for all $\gamma \in \Gamma$, we may consider the function $\Phi \in W(k)$ such that $\tilde{\Phi} = f$. This function is then left Γ -invariant. We will see in the next subsection that if $f \in N_k^p(\Gamma)$, then Φ is an automorphic form.

3. THE STRUCTURE THEOREM FOR CUSP FORMS

Let $\Gamma \subset \mathrm{SL}_2(\mathbb{Q})$ be a congruence subgroup. Let $\mathcal{A}(\Gamma)$ be the space of automorphic forms on $\mathrm{SL}_2(\mathbb{R})$, and let $\mathcal{A}(\Gamma)^\circ$ be the subspace of cusp forms. Recall that automorphic forms are required to be smooth, left Γ -invariant, K -finite, \mathcal{Z} -finite, and slowly increasing; we refer to [1] for the precise definitions. The spaces $\mathcal{A}(\Gamma)$ and $\mathcal{A}(\Gamma)^\circ$ are (\mathfrak{g}, K) -modules with respect to right translation. Let $\mathcal{A}_k(\Gamma)$ (resp. $\mathcal{A}_k(\Gamma)^\circ$) be the space of automorphic forms (resp. cusp forms) Φ satisfying $H \cdot \Phi = k\Phi$, or equivalently, $\Phi(gr(\theta)) = e^{ik\theta}\Phi(g)$ for all $\theta \in \mathbb{R}$ and $g \in \mathrm{SL}_2(\mathbb{R})$.

If $f \in \mathcal{A}(\Gamma)$ and $g \in \mathcal{A}(\Gamma)^\circ$, then the function $|fg|$ is integrable over $\Gamma \backslash \mathrm{SL}_2(\mathbb{R})$. In particular, $\mathcal{A}(\Gamma)^\circ \subset L^2(\Gamma \backslash \mathrm{SL}_2(\mathbb{R}))$. With respect to the L^2 inner product, the space $\mathcal{A}(\Gamma)^\circ$ decomposes into an orthogonal direct sum of irreducible representations, each occurring with finite multiplicity.

Let $\Phi \in \mathcal{A}(\Gamma)$. We will say that Φ is \mathfrak{n} -finite if $L^v \Phi = 0$ for large enough v . Let $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ be the space of \mathfrak{n} -finite automorphic forms, and let $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ be the subspace of \mathfrak{n} -finite cusp forms. The following properties are easy to verify:

- $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ is a (\mathfrak{g}, K) -submodule of $\mathcal{A}(\Gamma)$.

- $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ is the direct sum of its weight spaces, i.e.: If $\Phi \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ and $\Phi = \Phi_1 + \dots + \Phi_m$ with $\Phi_i \in \mathcal{A}_{k_i}(\Gamma)$ for different weights k_i , then $\Phi_i \in \mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}$ for each i .

Analogous statements hold for cusp forms.

Lemma 3.1. *Let k be an integer, and p a non-negative integer. Let Γ be a congruence subgroup of $\mathrm{SL}_2(\mathbb{Q})$. Let $f \in N_k^p(\Gamma)$ be non-zero. Define a function Φ on $\mathrm{SL}_2(\mathbb{R})$ by $\Phi(g) = (f|_k g)(i)$. Then $\Phi \in \mathcal{A}_k(\Gamma)_{\mathfrak{n}\text{-fin}}$. If f is a cusp form, then $\Phi \in \mathcal{A}_k(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$.*

Proof. Evidently, Φ is smooth, left Γ -invariant and has weight k . Since $N_k^p(\Gamma)$ is finite-dimensional (Lemma 1.2) and Ω acts on $N_k^p(\Gamma)$ (Lemma 1.1), the function f is $\mathbb{C}[\Omega]$ -finite. Hence, by Lemma 2.1, the function Φ is \mathcal{Z} -finite. The holomorphy of f at the cusps implies that Φ is slowly increasing. This proves $\Phi \in \mathcal{A}_k(\Gamma)$. Cuspidality of f translates into cuspidality of Φ . To prove \mathfrak{n} -finiteness, observe that $L^v f = 0$ for large enough v by Lemma 1.1. Hence $L^v \Phi = 0$ for large enough v by Lemma 2.1 and Lemma 1.3. \square

The following result is sometimes called the ‘‘duality theorem’’; see Theorem 2.10 of [2].

Proposition 3.2. *As (\mathfrak{g}, K) -modules, we have*

$$\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ = \bigoplus_{\ell=1}^{\infty} n_\ell \mathcal{D}_{\ell-1,+}, \quad n_\ell = \dim S_\ell(\Gamma).$$

The lowest weight vectors in the isotypical component $n_\ell \mathcal{D}_{\ell-1,+}$ correspond to elements of $S_\ell(\Gamma)$ via the map $\Phi \mapsto \tilde{\Phi}$, where $(\tilde{\Phi}|_k g)(i) = \Phi(g)$ for $g \in \mathrm{SL}_2(\mathbb{R})$.

Proof. Since $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ is a (\mathfrak{g}, K) -submodule of $\mathcal{A}(\Gamma)^\circ$, it decomposes into an orthogonal direct sum of irreducible (\mathfrak{g}, K) -modules. None of the irreducible constituents can be of the form $\mathcal{D}_{p,-}$ or a principal series representation, since any non-zero vector in such a constituent would not be \mathfrak{n} -finite. Neither can $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ contain any finite-dimensional representations; the lowest weight vector in such a constituent would give rise to a holomorphic cusp form of non-positive weight, which is not possible by Lemma 1.3. It follows that $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ can only contain constituents of the form $\mathcal{D}_{\ell-1,+}$ for $\ell \geq 1$. The fact that $\mathcal{D}_{\ell-1,+}$ occurs with multiplicity $\dim S_\ell(\Gamma)$ follows because a lowest weight vector in a constituent of the form $\mathcal{D}_{\ell-1,+}$ gives rise to an element of $S_\ell(\Gamma)$, and conversely. \square

Remark 3.3. It follows from Proposition 3.2 that $\mathcal{A}(\Gamma)_{\mathfrak{n}\text{-fin}}^\circ$ is an admissible (\mathfrak{g}, K) -module.

Knowing Proposition 3.2, it is now easy to derive the following Structure Theorem for cuspidal nearly holomorphic modular forms.

Theorem 3.4 (Structure theorem for cusp forms). *Fix non-negative integers k, p and a congruence subgroup Γ of $\mathrm{SL}_2(\mathbb{Q})$. There is an orthogonal direct sum decomposition*

$$N_k^p(\Gamma)^\circ = \bigoplus_{\substack{\ell \geq 1 \\ \ell \equiv k \pmod{2} \\ k-2p \leq \ell \leq k}} R^{(k-\ell)/2}(S_\ell(\Gamma)). \quad (9)$$

In particular, $N_0^p(\Gamma)^\circ = 0$ and $N_1^p(\Gamma)^\circ = S_1(\Gamma)$.

Proof. Let $f \in N_k^p(\Gamma)$. Define a function Φ on $\mathrm{SL}_2(\mathbb{R})$ by $\Phi(g) = (f|_k g)(i)$. Then $\Phi \in \mathcal{A}_k(\Gamma)_{\mathrm{n-fin}}^\circ$ by Lemma 3.1. If f has weight 0, then $f = 0$, since the weight 0 does not occur in $\mathcal{A}_k(\Gamma)_{\mathrm{n-fin}}^\circ$ by Proposition 3.2. Assume in the following that $k \geq 1$ and that f is non-zero.

Write $\Phi = \sum \Phi_i$, where each $\Phi_i \in \mathcal{A}_k(\Gamma)_{\mathrm{n-fin}}^\circ$ generates an irreducible (\mathfrak{g}, K) -module $V_i \cong \mathcal{D}_{\ell_i-1,+}$ with $\ell_i \geq 1$; this is possible by Proposition 3.2. Evidently, $f = \sum \tilde{\Phi}_i$, where $\tilde{\Phi}_i$ is the function on \mathbb{H}_1 corresponding to Φ_i via (7).

Since $f \in N_k^p(\Gamma)^\circ$, it follows from (2) of Lemma 1.1 that $L^{p+1}f = 0$, thus $L^{p+1}\Phi = 0$ by Lemma 2.1, and then also $L^{p+1}\Phi_i = 0$ for all i . The weight of $L^{p+1}\Phi_i$ being $k-2p-2$, it follows that V_i only contains weights greater or equal to $k-2p$. Hence $\ell_i \geq k-2p$ for all i .

Let $\Phi_{i,0} \in V_i$ be a lowest weight vector; thus $\Phi_{i,0}$ has weight $\ell_i \leq k$, and $\ell_i \equiv k \pmod{2}$. The corresponding function $\tilde{\Phi}_{i,0}$ on \mathbb{H}_1 is an element of $S_{\ell_i}(\Gamma)$. Since every weight occurs only once in V_i , we have $R^{(k-\ell_i)/2}\Phi_{i,0} = c_i\Phi_i$ for some non-zero constant c_i . By Lemma 2.1, it follows that

$$R^{(k-\ell_i)/2}\tilde{\Phi}_{i,0} = c_i\tilde{\Phi}_i,$$

and hence

$$f = \sum \tilde{\Phi}_i = \sum c_i^{-1}R^{(k-\ell_i)/2}\tilde{\Phi}_{i,0} \in \sum_{\substack{\ell \geq 1 \\ \ell \equiv k \pmod{2} \\ k-2p \leq \ell \leq k}} R^{(k-\ell)/2}(S_\ell(\Gamma)).$$

This proves that the left hand side of (9) is contained in the right hand side. The orthogonality of the right hand side follows from the above construction and the fact that the isotypical components in Proposition 3.2 are orthogonal; observe that the map $\Phi \mapsto \tilde{\Phi}$ is isometric with respect to the L^2 -scalar product on the left hand side and the Petersson inner product on the right hand side. \square

Remark 3.5. It is well known, or follows from an easy calculation, that Ω acts on $\mathcal{D}_{\ell-1,+}$ by the scalar $\frac{1}{2}\ell(\frac{1}{2}\ell-1)$. Hence, by Lemma 2.1, Ω acts on the subspace $R^{(k-\ell)/2}(S_\ell(\Gamma))$ of $N_k^p(\Gamma)^\circ$ by $\frac{1}{2}\ell(\frac{1}{2}\ell-1)$. In particular, Ω acts diagonalizably on $N_k^p(\Gamma)^\circ$, and the pieces in the decomposition (9) can be intrinsically characterized as the eigenspaces with respect to Ω .

Petersson inner products. For $f, g \in N_k(\Gamma)$ with at least one of them in $N_k(\Gamma)^\circ$, we define the Petersson inner product $\langle f, g \rangle$ by the equation

$$\langle f, g \rangle = \mathrm{vol}(\Gamma \backslash \mathbb{H}_1)^{-1} \int_{\Gamma \backslash \mathbb{H}_1} f(\tau) \overline{g(\tau)} \frac{dx dy}{y^2}.$$

It can be easily checked that

$$\langle f, g \rangle = \langle \Phi_f, \Phi_g \rangle, \quad (10)$$

where $\Phi_f(h) = (f|_k h)(i)$ (and Φ_g is defined similarly) and the inner product of Φ_f and Φ_g is defined by

$$\langle \Phi_f, \Phi_g \rangle = \frac{1}{\mathrm{vol}(\mathrm{SL}_2(\mathbb{Z}) \backslash \mathrm{SL}_2(\mathbb{R}))} \int_{\mathrm{SL}_2(\mathbb{Z}) \backslash \mathrm{SL}_2(\mathbb{R})} \Phi_f(h) \overline{\Phi_g(h)} dh.$$

We define the subspace $\mathcal{E}_k(\Gamma)$ to be the orthogonal complement of $N_k(\Gamma)^\circ$ in $N_k(\Gamma)$. Let $\mathcal{E}_k^p(\Gamma) = \mathcal{E}_k(\Gamma) \cap N_k^p(\Gamma)$. We write $E_k(\Gamma)$ to mean $\mathcal{E}_k^0(\Gamma)$. In Corollary 4.4 below we will prove that $N_k^p(\Gamma)$ is the orthogonal direct sum of $N_k^p(\Gamma)^\circ$ and $\mathcal{E}_k^p(\Gamma)$.

Lemma 3.6. *Let k be a non-negative integer. Let $f \in \mathcal{E}_k(\Gamma)$, and let $\Phi_f \in \mathcal{A}(\Gamma)_{\text{n-fin}}$ be the corresponding function on $\text{SL}_2(\mathbb{R})$. Then Φ_f is orthogonal to $\mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$.*

Proof. Let $\Psi \in \mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$; we have to show that $\langle \Phi_f, \Psi \rangle = 0$. We may assume that Ψ generates an irreducible module $\mathcal{D}_{\ell-1,+}$ for some $\ell \geq 1$. Since Φ_f has weight k , we may assume that Ψ does as well. But then Ψ corresponds to an element g of $N_k(\Gamma)^\circ$. By hypothesis $\langle f, g \rangle = 0$. Hence $\langle \Phi_f, \Psi \rangle = 0$ by (10). \square

Lemma 3.7. *Let $k \geq 1$ and $v \geq 0$ be integers. Then R^v takes $S_k(\Gamma)$ to $N_{k+2v}^v(\Gamma)^\circ$ and $E_k(\Gamma)$ to $\mathcal{E}_{k+2v}^v(\Gamma)$.*

Proof. The fact that R^v takes $S_k(\Gamma)$ to $N_{k+2v}^v(\Gamma)^\circ$ is an immediate consequence of the fact that the differential operator R commutes with the $|_k$ operator and does not increase the support of the Fourier coefficients.

Let us show that R^v takes $E_k(\Gamma)$ to $\mathcal{E}_{k+2v}^v(\Gamma)$. Let $f \in E_k(\Gamma)$. In view of (10), it suffices to show that $R^v(\Phi_f)$ and Φ_g are orthogonal for all $g \in N_{k+2v}^v(\Gamma)^\circ$. But note that $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})\Phi_f$ and $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})\Phi_g$ are orthogonal submodules of $\mathcal{A}(\Gamma)$ (as $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})\Phi_g$ is completely contained in $\mathcal{A}(\Gamma)^\circ$ and $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})\Phi_f$ is contained in the orthogonal complement of $\mathcal{A}(\Gamma)^\circ$ by Lemma 3.6). Hence $R^v(\Phi_f)$ and Φ_g are orthogonal. \square

Lemma 3.8. *Let $f \in S_k(\Gamma)$. Then for all $v \geq 0$, there exists a constant $c_{k,v}$ (depending only on k, v) such that*

$$\langle R^v(f), R^v(f) \rangle = c_{k,v} \langle f, f \rangle.$$

Proof. Consider the (\mathfrak{g}, K) module $\mathcal{D}_{k-1,+}$ and let v_0 be a lowest-weight vector in it. Note that v_0 is unique up to multiples. It is well-known that $\mathcal{D}_{k-1,+}$ is unitarizable; let $\langle \cdot, \cdot \rangle$ denote the (unique up to multiples) \mathfrak{g} -invariant inner product on it. Put $c_{k,v} = \langle R^v(v_0), R^v(v_0) \rangle / \langle v_0, v_0 \rangle$. Note that $c_{k,v}$ does not depend on the choice of model for $\mathcal{D}_{k-1,+}$, the choice of v_0 or the normalization of inner products.

Now all we need to observe is that the automorphic form $\Phi_f \in \mathcal{A}(\Gamma)_{\text{n-fin}}^\circ$ corresponding to f generates a module isomorphic to $\mathcal{D}_{k-1,+}$, that Φ_f is a lowest weight vector in this module, and (10). \square

Proposition 3.9. *Let $f \in M_k(\Gamma)$, $g \in S_k(\Gamma)$. Then*

$$\langle R^v(f), R^v(g) \rangle = c_{k,v} \langle f, g \rangle,$$

where the constant $c_{k,v}$ is as in the previous lemma.

Proof. Because of Lemma 3.7, we may assume that f and g both belong to $S_k(\Gamma)$. Now the Proposition follows by applying the previous lemma to $f + g$. \square

4. THE NON-CUSPIDAL CASE

The obstruction in the non-cuspidal case. The Structure Theorem 3.4 cannot hold without modifications for non-cuspidal nearly holomorphic modular forms. The reason is the existence of the weight 2 Eisenstein series

$$E_2(\tau) = -\frac{3}{\pi y} + 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) e^{2\pi i n \tau}, \quad \sigma_1(n) = \sum_{d|n} d. \quad (11)$$

As is well known, E_2 is modular with respect to $\text{SL}_2(\mathbb{Z})$; thus, $E_2 \in N_2^1(\Gamma)$ for any congruence subgroup Γ of $\text{SL}_2(\mathbb{Z})$. But evidently E_2 cannot be obtained via raising operators from holomorphic forms of lower weight, since the only modular forms of weight 0 are the constant functions.

Let $\Phi_2 \in \mathcal{A}_2(\Gamma)_{\text{n-fin}}$ be the automorphic form corresponding to E_2 via Lemma 3.1. Let V_{Φ_2} be the (\mathfrak{g}, K) -module generated by Φ_2 . Since $L_2\Phi_2 = \frac{3}{\pi}$, we have $L\Phi_2 \in \mathbb{C}$ by Lemma 2.1. The weight structure of V_{Φ_2} is therefore $[0, 2, 4, \dots]$, and the constant functions are a submodule of V_{Φ_2} . More precisely, there is an exact sequence

$$0 \longrightarrow \mathbb{C} \longrightarrow V_{\Phi_2} \longrightarrow \mathcal{D}_{1,+} \longrightarrow 0; \quad (12)$$

recall that $\mathcal{D}_{1,+}$ is the lowest weight module with weight structure $[2, 4, 6, \dots]$. Clearly, this sequence does not split. Consequently, unlike in the cuspidal case, the space $\mathcal{A}(\Gamma)_{\text{n-fin}}$ is not the direct of irreducible submodules. However, the following result states that V_{Φ_2} represents the only obstruction:

Proposition 4.1. *As (\mathfrak{g}, K) -modules, we have*

$$\mathcal{A}(\Gamma)_{\text{n-fin}} = V_{\Phi_2} \oplus \bigoplus_{\ell=1}^{\infty} n_{\ell} \mathcal{D}_{\ell-1,+}, \quad n_{\ell} = \dim M_{\ell}(\Gamma).$$

The lowest weight vectors in the isotypical component $n_{\ell} \mathcal{D}_{\ell-1,+}$ correspond to elements of $M_{\ell}(\Gamma)$ via the map $\Phi \mapsto \tilde{\Phi}$, where $(\tilde{\Phi}|_k g)(i) = \Phi(g)$ for $g \in \text{SL}_2(\mathbb{R})$. The module V_{Φ_2} sits in the exact sequence (12) and is generated by the function Φ_2 such that $\tilde{\Phi}_2 = E_2$.

To prove this result, we will set up a certain algebraic apparatus. It turns out that the mechanism of category \mathcal{O} is well suited toward our problem. In the SL_2 case this mechanism could be replaced by more direct arguments, but we prefer to use category \mathcal{O} because this method generalizes to the Sp_4 case; see [6]. Our reference for category \mathcal{O} will be [3].

Roots and weights. Let H, R, L be the elements of $\mathfrak{g}_{\mathbb{C}}$ defined in (5). Then $\mathfrak{h} = \langle H \rangle$ is a Cartan subalgebra of $\mathfrak{g}_{\mathbb{C}}$. Let $\Phi \subset \mathfrak{h}^*$ be the root system of $\mathfrak{g}_{\mathbb{C}}$ corresponding to \mathfrak{h} . Then R and L span the corresponding root spaces. We identify an element λ of \mathfrak{h}^* with the element $\lambda(H)$ of \mathbb{C} . Then $\Phi = \{\pm 2\}$. Let E be the \mathbb{R} -span of Φ . We endow E with the inner product (\cdot, \cdot) given by the usual multiplication of real numbers. Perhaps counterintuitively, we will declare -2 to be a positive (and simple) root, with corresponding root vector L , and $+2$ a negative root, with corresponding root vector R . The weight lattice Λ is defined as

$$\Lambda = \left\{ \lambda \in E \mid 2 \frac{(\lambda, \alpha)}{(\alpha, \alpha)} \in \mathbb{Z} \text{ for all } \alpha \in \Phi \right\}. \quad (13)$$

Evidently, $\Lambda = \mathbb{Z} \subset E$. There is an ordering on Λ defined as follows:

$$\mu \preceq \lambda \iff \lambda \in \mu + \Gamma, \quad (14)$$

where Γ is the set of all $\mathbb{Z}_{\geq 0}$ -multiples of the positive root. Hence,

$$\mu \preceq \lambda \iff \lambda \leq \mu \text{ and } \lambda \equiv \mu \pmod{2}.$$

The fundamental weight is -1 , and the dominant integral weights are the $\mathbb{Z}_{\geq 0}$ -multiples of the fundamental weight. We write Λ^+ for the set of dominant integral weights. Hence, $\Lambda^+ = \{-1, -2, \dots\}$. We write $\varrho = -1$ for half the sum of the positive roots. As before, let \mathcal{Z} denote the center of the universal enveloping algebra $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$. Via the Harish-Chandra isomorphism, all possible characters of \mathcal{Z} are indexed by elements of \mathfrak{h}^* modulo Weyl group action; see equation (8.32) in [4]. We denote by χ_{λ} the character of \mathcal{Z} corresponding to $\lambda \in \mathfrak{h}^*$. Note that $\chi_{w\lambda} = \chi_{\lambda}$ for all $w \in W$, where the Weyl group acts by negation on $\mathfrak{h}^* \cong \mathbb{C}$.

Verma modules. We recall the definition of the standard Verma modules. Let λ be an integer, considered as an element of the weight lattice Λ . Let \mathbb{C}_λ be the one-dimensional space on which $\mathfrak{h} = \langle H \rangle$ acts via λ . Let

$$\mathfrak{b} = \mathfrak{h} + \langle L \rangle$$

be the Borel algebra defined by our positive system. We consider \mathbb{C}_λ a \mathfrak{b} -module with the action of L being trivial. Then the *Verma module* corresponding to λ is defined as

$$N(\lambda) = \mathcal{U}(\mathfrak{g}_{\mathbb{C}}) \otimes_{\mathcal{U}(\mathfrak{b})} \mathbb{C}_\lambda. \quad (15)$$

Clearly, $N(\lambda)$ contains the weight λ with multiplicity one. Any non-zero vector v in $N(\lambda)$ of weight λ is called a *highest weight vector*. It is well known that $N(\lambda)$ has the following properties:

- (1) $N(\lambda)$ is a free module of rank 1 over $\mathcal{U}(R) = \mathbb{C}[R]$.
- (2) The set of weights of $N(\lambda)$ is $\lambda - \Gamma = \{\lambda, \lambda + 2, \dots\}$. Each weight occurs with multiplicity one.
- (3) The module $N(\lambda)$ is a universal highest weight module for the weight λ , meaning it satisfies this universal property: Let M be a $\mathfrak{g}_{\mathbb{C}}$ -module which contains a vector v with the following properties:
 - $M = \mathcal{U}(\mathfrak{g}_{\mathbb{C}})v$;
 - v has weight λ ;
 - $Lv = 0$.

Then there exists a surjection $N(\lambda) \rightarrow M$ mapping a highest weight vector in $N(\lambda)$ to v .

- (4) $N(\lambda)$ admits a unique irreducible submodule, and a unique irreducible quotient $L(\lambda)$. In particular, $N(\lambda)$ is indecomposable. See Theorem 1.2 of [3].
- (5) $N(\lambda)$ has finite length. Each factor in a composition series is of the form $L(\mu)$ for some $\mu \leq \lambda$.
- (6) $N(\lambda)$ admits the central character $\chi_{\lambda+\varrho}$, i.e., \mathcal{Z} acts on $N(\lambda)$ via $\chi_{\lambda+\varrho}$. See Sects. 1.7–1.10 of [3]. Note that Humphrey's χ_λ is our $\chi_{\lambda+\varrho}$.
- (7) $L(\lambda)$ is finite-dimensional if and only if $\lambda \in \Lambda^+ = \{0, -1, -2, \dots\}$. See Theorem 1.6 of [3].
- (8) $N(\lambda)$ is simple if and only if $\lambda > 0$. See Theorem 4.4 of [3].

Evidently, $L(0)$ is the trivial $\mathfrak{g}_{\mathbb{C}}$ -module. It has central character χ_ϱ .

Category \mathcal{O} . We recall from Sect. 1.1 of [3] the definition of category \mathcal{O} . This category is defined with respect to a choice of Cartan subalgebra \mathfrak{h} and a choice of simple roots, and we make the choices specified above. Let \mathfrak{n} be the space spanned by the positive root vectors, hence, in our case, $\mathfrak{n} = \langle L \rangle$. A $\mathfrak{g}_{\mathbb{C}}$ -module M is said to be in category \mathcal{O} if it satisfies the following conditions:

- ($\mathcal{O}1$) M is a finitely generated $\mathcal{U}(\mathfrak{g}_{\mathbb{C}})$ -module.
- ($\mathcal{O}2$) M is the direct sum of its weight spaces, and all weights are integral.
- ($\mathcal{O}3$) M is locally \mathfrak{n} -finite. This means: For each $v \in M$ the subspace $\mathcal{U}(\mathfrak{n})v$ is finite-dimensional.

Note that we are slightly varying the definition of category \mathcal{O} by requiring that all weights are integral; the relevant results in [3] still hold with this modification.

\mathcal{O} is an abelian category. Evidently, \mathcal{O} contains all Verma modules $N(\lambda)$ and their irreducible quotients $L(\lambda)$. The modules M in \mathcal{O} have many nice properties, as explained in the first sections of [3]. For example:

- M has finite length, and admits a filtration

$$0 = V_0 \subset V_1 \subset \dots \subset V_n \subset M, \quad (16)$$

with $V_i/V_{i-1} \cong L(\lambda)$ for some $\lambda \in \mathfrak{h}^*$.

- M can be written as a finite direct sum of indecomposable modules.
- If M is an indecomposable module, then there exists a character χ of \mathcal{Z} such that $M = N(\chi)$. Here,

$$N(\chi) = \{v \in M \mid (z - \chi(z))^n v = 0 \text{ for some } n \text{ depending on } z\}. \quad (17)$$

For each M in category \mathcal{O} we may write

$$M = \bigoplus_{\chi} N(\chi), \quad (18)$$

where χ runs over characters of \mathcal{Z} , and $N(\chi)$ is defined as in (17); see Sect. 1.12 of [3]. The *modules* $N(\chi)$ may or may not be indecomposable.

Another feature of \mathcal{O} is the existence of a duality functor $M \mapsto M^\vee$, as explained in Sect. 3.2 of [3]. In general, M^\vee is *not* the contragredient of M , as \mathcal{O} is not closed under taking contragredients. The duality functor in \mathcal{O} has the following properties:

- $M \mapsto M^\vee$ is exact and contravariant.
- $M^{\vee\vee} \cong M$.
- $(M_\chi)^\vee \cong (M^\vee)_\chi$ for a character χ of \mathcal{Z} .
- $L(\lambda)^\vee \cong L(\lambda)$.
- $\text{Ext}_{\mathcal{O}}(M, N) \cong \text{Ext}_{\mathcal{O}}(N^\vee, M^\vee)$. See Sect. 3.1 of [3] for the definition of the $\text{Ext}_{\mathcal{O}}$ groups.

Evidently, $L(0)$ is the trivial $\mathfrak{g}_{\mathbb{C}}$ -module. It is easy to see that there is an exact sequence

$$0 \longrightarrow L(2) \longrightarrow N(0) \longrightarrow L(0) \longrightarrow 0. \quad (19)$$

Since $N(0)$ is indecomposable, this sequence does not split. Applying the duality functor, we get another non-split exact sequence

$$0 \longrightarrow L(0) \longrightarrow N(0)^\vee \longrightarrow L(2) \longrightarrow 0. \quad (20)$$

It is an exercise to show that the sequence (19) is the only non-trivial extension of $L(0)$ by $L(2)$; similarly for (20). The fact that $\text{Ext}_{\mathcal{O}}(L(0), L(2)) = 1$ can also be seen by applying the functor $\text{Hom}_{\mathcal{O}}(_, L(2))$ to (19) and considering the resulting long exact sequence.

Proof of Proposition 4.1. Before starting the proof of Proposition 4.1, let us comment on the relationship between (\mathfrak{g}, K) -modules and $\mathfrak{g}_{\mathbb{C}}$ -modules. Clearly, every (\mathfrak{g}, K) -module is also a $\mathfrak{g}_{\mathbb{C}}$ -module. Conversely, let (π, V) be a $\mathfrak{g}_{\mathbb{C}}$ -module all of whose weights are integral, and such that V is the direct sum of its weight spaces. If $v \in V$ has weight k , i.e., if $\pi(H)v = kv$, then we define a K -action on $\mathbb{C}v$ by $\pi(r(\theta))v = e^{ik\theta}v$. Since V is the direct sum of its weight spaces, this defines a K -action on all of V . One can verify that, with this K -action, V becomes a (\mathfrak{g}, K) -module. In particular, every module in category \mathcal{O} is naturally a (\mathfrak{g}, K) -module. The upshot is that in the following arguments we do not have to worry about the distinction between (\mathfrak{g}, K) -modules and $\mathfrak{g}_{\mathbb{C}}$ -modules.

With these comments in mind, it is clear that we have the following isomorphisms of (\mathfrak{g}, K) -modules:

- $\mathcal{F}_p \cong L(-p + 1)$ for $p \geq 1$;
- $\mathcal{D}_{p,+} \cong L(p + 1)$ for $p \geq 0$;
- $V_{\Phi_2} \cong N(0)^\vee$.

The third isomorphism follows by comparing the exact sequences (12) and (20), observing the uniqueness comment made after (20).

We are now ready to prove Proposition 4.1. As we saw, the modular form E_2 gives rise to the submodule V_{Φ_2} of $\mathcal{A}(\Gamma)_{\text{n-fin}}$. Every non-zero $f \in M_\ell(\Gamma)$ gives rise to a copy of $\mathcal{D}_{\ell-1,+}$ inside $\mathcal{A}(\Gamma)_{\text{n-fin}}$. It is therefore clear that

$$\mathcal{A}(\Gamma)_{\text{n-fin}} \supset V_{\Phi_2} \oplus \bigoplus_{\ell=1}^{\infty} n_\ell \mathcal{D}_{\ell-1,+}, \quad n_\ell = \dim M_\ell(\Gamma). \quad (21)$$

To prove the converse, observe first that

- $\mathcal{A}(\Gamma)_{\text{n-fin}}$ contains no negative weights;
- $\mathcal{A}(\Gamma)_{\text{n-fin}}$ contains the weight 0 exactly once, the corresponding weight space consisting of the constant functions.

Both statements follow from Lemma 1.3. We define

$$\mathcal{A}_{\leq k}(\Gamma)_{\text{n-fin}} = \sum_{\ell=0}^k \mathcal{A}_\ell(\Gamma)_{\text{n-fin}},$$

and let $\mathcal{A}_{\langle \leq k \rangle}(\Gamma)_{\text{n-fin}}$ be the (\mathfrak{g}, K) -module generated by elements of $\mathcal{A}_{\leq k}(\Gamma)_{\text{n-fin}}$. Evidently,

$$\mathcal{A}_{\langle \leq k \rangle}(\Gamma)_{\text{n-fin}} \supset V_{\Phi_2} \oplus \bigoplus_{\ell=1}^k n_\ell \mathcal{D}_{\ell-1,+}, \quad n_\ell = \dim M_\ell(\Gamma). \quad (22)$$

To prove equality in (21), it is enough to prove equality in (22).

It follows from the finite-dimensionality of the spaces $M_\ell(\Gamma)$ that $\mathcal{A}_{\leq k}(\Gamma)_{\text{n-fin}}$ is finite-dimensional. Hence $\mathcal{A}_{\langle \leq k \rangle}(\Gamma)_{\text{n-fin}}$ is finitely generated. This proves that $\mathcal{A}_{\langle \leq k \rangle}(\Gamma)_{\text{n-fin}}$ is in category \mathcal{O} . By properties of this category, we may write

$$\mathcal{A}_{\langle \leq k \rangle}(\Gamma)_{\text{n-fin}} = V_1 \oplus \dots \oplus V_n$$

with indecomposable submodules. Let

$$0 = V_{i,0} \subset V_{i,1} \subset \dots \subset V_{i,n_i} = V_i \quad (23)$$

be a filtration for V_i such that $V_{i,j}/V_{i,j-1} \cong L(\lambda_{i,j})$ for some $\lambda_{i,j} \in \mathbb{Z}$. Since there are no negative weights, we have $\lambda_{i,j} \geq 0$ for all i and j .

Let χ_i be the character of \mathcal{Z} such that $V_i = V_i(\chi_i)$; see (17). We think of χ_i as a non-negative integer. Assume that $\chi_i > 1$ or $\chi_i = 0$. Then $\lambda_{i,j} = \chi_i + 1$ for all j , since, among the $L(\lambda)$ with $\lambda \geq 0$, only $L(\chi_i + 1)$ has central character χ_i . Now $\text{Ext}_{\mathcal{O}}(L(\lambda), L(\lambda)) = 0$ for all λ by Proposition 3.1 d) of [3]. It follows that V_i is a direct sum of copies of $L(\lambda_i)$, where $\lambda_i := \chi_i + 1$. Since V_i is indecomposable, we must have $V_i = L(\lambda_i)$.

Now consider a V_i with $\chi_i = 1$ (i.e., $\chi_i = \chi_\varrho$). The only $L(\lambda)$ with $\lambda \geq 0$ and this central character are $L(0)$ and $L(2)$. If $L(0)$ does not occur in V_i , then the same argument as above applies, and we see that $V_i = L(2)$. Assume that $L(0)$ does occur in V_i . Since the weight 0 occurs exactly once in the entire space, there is exactly one V_i with this property, and this V_i contains $L(0)$ exactly once. Since the weight 0 space consists of the constant functions, it appears as a subrepresentation in V_i . Hence, we may assume it occurs at the bottom of the filtration, i.e., $V_{i,1} = L(0)$. If V_i would not contain any $L(2)$ subquotients, then $\mathcal{A}_{\langle \leq k \rangle}(\Gamma)_{\text{n-fin}}$ would be completely reducible, which we know is not the case. Hence there is at least one $L(2)$ subquotient sitting on top of the $L(0)$. By (20) and the remark following it, $V_{i,2} \cong N(0)^\vee$. Now

$$\text{Ext}_{\mathcal{O}}(L(2), N(0)^\vee) \cong \text{Ext}_{\mathcal{O}}(N(0), L(2)) = 0$$

by Proposition 1.3 b) of [3]. This means that there can be no further $L(2)$'s on top of the $N(0)^\vee$, and it follows that $V_i = N(0)^\vee$.

To summarize, we proved that, abstractly,

$$\mathcal{A}_{(\leq k)}(\Gamma)_{\text{n-fin}} = N(0)^\vee \oplus \bigoplus_{\lambda=1}^{\infty} m_\lambda L(\lambda)$$

with non-negative integers m_λ almost all of which are zero. A moment's consideration shows that $m_\lambda = 0$ for $\lambda > k$. Since $L(\lambda) \cong \mathcal{D}_{\lambda-1,+}$ and $N(0)^\vee \cong V_{\Phi_2}$, and since we know $\mathcal{D}_{\lambda-1,+}$ cannot occur more than $\dim M_\lambda(\Gamma)$ times, comparison with (22) shows that we must have equality in (22). This concludes the proof of Proposition 4.1. \square

The Structure Theorem for all modular forms. We can now provide an alternative proof of Theorem 5.2 of [7].

Theorem 4.2 (Structure theorem for all modular forms). *Fix non-negative integers k, p and a congruence subgroup Γ of $\text{SL}_2(\mathbb{Z})$. Then:*

- (1) $N_0^p(\Gamma) = \mathbb{C}$.
- (2) If k is even and $2 \leq k < 2 + 2p$, then

$$N_k^p(\Gamma) = R^{(k-2)/2}(\mathbb{C}E_2) \oplus \bigoplus_{\substack{\ell \geq 1 \\ \ell \equiv k \pmod{2} \\ k-2p \leq \ell \leq k}} R^{(k-\ell)/2}(M_\ell(\Gamma)). \quad (24)$$

- (3) If k is odd, or if k is even and $k \geq 2 + 2p$, then

$$N_k^p(\Gamma) = \bigoplus_{\substack{\ell \geq 1 \\ \ell \equiv k \pmod{2} \\ k-2p \leq \ell \leq k}} R^{(k-\ell)/2}(M_\ell(\Gamma)). \quad (25)$$

Proof. The proof is analogous to that of Theorem 3.4. Instead of Proposition 3.2, we use Proposition 4.1. Observe that $R^{(k-2)/2}E_2$ is in $N_k^{k/2}(\Gamma)$, but not in $N_k^{k/2-1}(\Gamma)$, so in order for E_2 to contribute to $N_k^p(\Gamma)$ (for k even) we must have $\frac{k}{2} \leq p$, or equivalently, $k < 2 + 2p$. \square

A simplified version of the Structure Theorem for all modular forms would be this: If $p < \frac{k-2}{2}$, then

$$N_k^p(\Gamma) = \bigoplus_{\substack{\ell \geq 1 \\ \ell \equiv k \pmod{2} \\ k-2p \leq \ell \leq k}} R^{(k-\ell)/2}(M_\ell(\Gamma)). \quad (26)$$

The hypothesis $p < \frac{k-2}{2}$ implies that, in the arguments in the proof of the theorem, we never “reach down” to weight 2. Hence, the component V_{Φ_2} appearing in Proposition 4.1 can be ignored.

Corollary 4.3 (Structure theorem for non-cusp forms). *Fix non-negative integers k, p and a congruence subgroup Γ of $\text{SL}_2(\mathbb{Z})$. Then:*

- (1) $\mathcal{E}_0^p(\Gamma) = \mathbb{C}$.
- (2) If k is even and $2 \leq k < 2 + 2p$, then

$$\mathcal{E}_k^p(\Gamma) = R^{(k-2)/2}(\mathbb{C}E_2) \oplus \bigoplus_{\substack{\ell \geq 1 \\ \ell \equiv k \pmod{2} \\ k-2p \leq \ell \leq k}} R^{(k-\ell)/2}(E_\ell(\Gamma)). \quad (27)$$

- (3) If k is odd, or if k is even and $k \geq 2 + 2p$, then

$$\mathcal{E}_k^p(\Gamma) = \bigoplus_{\substack{\ell \geq 1 \\ \ell \equiv k \pmod{2} \\ k-2p \leq \ell \leq k}} R^{(k-\ell)/2}(E_\ell(\Gamma)). \quad (28)$$

Proof. That the right side of each equation is contained in the left side is immediate from Lemma 3.7 and the fact that Φ_2 (the automorphic form corresponding to E_2) lies in the orthogonal complement of the cusp forms. That the left side is contained in the right side follows from Theorem 4.2, the fact that $M_\ell(\Gamma)$ is the orthogonal sum of $S_\ell(\Gamma)$ and $E_\ell(\Gamma)$, and the fact that the R^v maps preserve inner products up to a constant. \square

Corollary 4.4. *Let k and p be non-negative integers. The space $N_k^p(\Gamma)$ is the orthogonal direct sum of $N_k^p(\Gamma)^\circ$ and $\mathcal{E}_k^p(\Gamma)$.*

Proof. By the structure theorems, it is enough to prove the assertion for $p = 0$. In this case the claim is that $M_k(\Gamma)$ is the orthogonal direct sum of $S_k(\Gamma)$ and $E_k(\Gamma)$. Clearly,

$$M_k(\Gamma) = S_k(\Gamma) \oplus S_k(\Gamma)^\perp \quad \text{and} \quad E_k(\Gamma) \subset S_k(\Gamma)^\perp.$$

Hence, our task is to show that a non-zero element f of $S_k(\Gamma)^\perp$ is orthogonal to all of $N_k(\Gamma)^\circ$. Let Φ_f be the function on $\mathrm{SL}_2(\mathbb{R})$ corresponding to f . We will in fact show that Φ_f is orthogonal to any cusp form Ψ . We may assume that Ψ generates an irreducible representation $\mathcal{D}_{\ell-1,+}$. Assume first that Ψ has weight ℓ , i.e., Ψ is the lowest weight vector in $\mathcal{D}_{\ell-1,+}$. If $\ell \neq k$, then $\langle \Phi, \Psi \rangle = 0$ since the weights do not match. If $\ell = k$, then $\langle \Phi, \Psi \rangle = 0$ since Ψ corresponds to an element of $S_k(\Gamma)$. Now assume that Ψ has weight greater than ℓ . Then $\Psi = R\Psi'$ for some $\Psi' \in \mathcal{D}_{\ell-1,+}$, and the general formula

$$\langle \Phi, R\Psi' \rangle + \langle L\Phi, \Psi' \rangle = 0$$

shows that $\langle \Phi, R\Psi' \rangle = 0$, because Φ is a lowest weight vector. This concludes the proof. \square

Remark 4.5. It is well-known that $E_k(\Gamma) = S_k(\Gamma)^\perp$ is spanned by the various weight k holomorphic Eisenstein series on Γ .

REFERENCES

- [1] Armand Borel and Hervé Jacquet. Automorphic forms and automorphic representations. In *Automorphic forms, representations and L-functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1*, Proc. Sympos. Pure Math., XXXIII, pages 189–207. Amer. Math. Soc., Providence, R.I., 1979. With a supplement “On the notion of an automorphic representation” by R. P. Langlands.
- [2] Stephen Gelbart. *Automorphic forms on adèle groups*. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1975. Annals of Mathematics Studies, No. 83.
- [3] James Humphreys. *Representations of semisimple Lie algebras in the BGG category \mathcal{O}* , volume 94 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2008.
- [4] Anthony Knapp. *Representation theory of semisimple groups*, volume 36 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1986. An overview based on examples.
- [5] Toshitsune Miyake. *Modular forms*. Springer-Verlag, Berlin, 1989. Translated from the Japanese by Yoshitaka Maeda.
- [6] Ameya Pitale, Abhishek Saha, and Ralf Schmidt. Lowest weight modules of $\mathrm{Sp}_4(\mathbb{R})$ and nearly holomorphic Siegel modular forms. Preprint, 2014.
- [7] Goro Shimura. Nearly holomorphic functions on Hermitian symmetric spaces. *Math. Ann.*, 278(1-4):1–28, 1987.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OK 73019, USA
E-mail address: apitale@math.ou.edu

DEPARTMENTS OF MATHEMATICS, UNIVERSITY OF BRISTOL, BRISTOL BS81SN, UK
E-mail address: abhishek.saha@bris.ac.uk

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF OKLAHOMA, NORMAN, OK 73019, USA
E-mail address: rschmidt@math.ou.edu