

**I.** Let  $f(x, y, z)$  be the function  $ye^{xz^2}$ .

(8)

1. Calculate the gradient  $\nabla f(x, y, z)$ .

2. Calculate the rate of change of  $f$  at the point  $(2, 1, 1)$ , in the direction toward the point  $(0, 0, 2)$ .

3. Let  $S$  be the level surface of  $f$  through the point  $(2, 1, 1)$ . Find an equation for the tangent plane to  $S$  at the point  $(2, 1, 1)$ .

**II.** The picture at the right shows a certain vector field  $P\vec{i} + Q\vec{j}$

(6) in the  $xy$ -plane.

1. Is  $\frac{\partial Q}{\partial x}$  positive, negative, or zero?


2. Is  $\frac{\partial Q}{\partial y}$  positive, negative, or zero?

3. Is  $\frac{\partial P}{\partial x}$  positive, negative, or zero?

4. Is  $\frac{\partial P}{\partial y}$  positive, negative, or zero?


5. If  $C$  is the line segment from  $(1, -1)$  to  $(1, 0)$ , is  $\int_C (P\vec{i} + Q\vec{j}) \cdot d\vec{r}$  positive, negative, or zero?

6. If  $C$  is the line segment from  $(1, 0)$  to  $(1, 1)$ , is  $\int_C (P\vec{i} + Q\vec{j}) \cdot d\vec{r}$  positive, negative, or zero?



- III.** Let  $C$  consist of the clockwise arc of the circle  $x^2 + y^2 = 4$  from  $(2, 0)$  to  $(0, 2)$ , followed by the line segment  
 (7) from  $(0, 2)$  to  $(2, 4)$ . Calculate  $\int_C (2xy \cos(x^2) \vec{i} + (\sin(x^2) + \cos(y)) \vec{j}) \cdot d\vec{r}$ .

- IV.** The following two questions refer to the function  $f(x, y)$  whose graph is shown here, and the four functions  
 (4) indicated in the graphs A, B, C, and D. All of the functions have domain the points  $(x, y)$  with  $0 \leq x \leq 1$  and  $0 \leq y \leq 1$ .



- The graph of the function  $\frac{\partial f}{\partial y}$  looks most like the function in  
 (a) A                      (b) B                      (c) C                      (d) D
- The graph of the function  $\frac{\partial f}{\partial x}$  looks most like the function in  
 (a) A                      (b) B                      (c) C                      (d) D

**V.** A curve  $C$  is parameterized as  $\vec{r}(t) = e^t \vec{i} + \ln(\cos(t)) \vec{j}$  for  $0 \leq t \leq 1$ . Using this parameterization, (6) express each of the following as an ordinary definite integral of a function of  $t$ . Make reasonable algebraic simplifications, but *do not* try to calculate the actual values of the integrals.

1.  $\int_C \ln(x) e^y ds$

2.  $\int_C e^x dy$

3.  $\int_C (x^2 \vec{i} + e^y \vec{j}) \cdot d\vec{r}$

**VI.** Rewrite the following integrals using polar or spherical coordinates. Be sure to supply the correct limits (8) of integration for the new variables. *Do not* carry out the calculation of the integrals.

1.  $\int_0^1 \int_{-\sqrt{1-x^2}}^0 e^{x^2+y^2} dy dx$

2.  $\int_0^3 \int_0^{\sqrt{9-x^2}} \int_0^{\sqrt{9-x^2-y^2}} z \sqrt{x^2 + y^2 + z^2} dz dy dx$

**VII.** Let  $r$  and  $\theta$  be the polar coordinates in the  $xy$ -plane. For the function  $f(x, y) = x^3y$ , calculate  $\frac{\partial f}{\partial \theta}$  in terms of  $x$  and  $y$ .

**VIII.** Let  $\vec{F}(x, y, z) = -xz\vec{i} - yz\vec{j} + z^2\vec{k}$ , and let the surface  $S$  be the ellipsoid  $\frac{x^2}{a^2} + \frac{y^2}{b^2} + \frac{z^2}{c^2} = 1$ . Calculate  $\iint_S \vec{F} \cdot d\vec{S}$ , where  $S$  has the outward normal.



**IX.** The picture at the right shows a parameterization of the outer surface  $S$  of a football. It is parameterized by letting  $\theta$  be the polar angle in the  $xz$ -plane, and  $h$  be the  $y$ -coordinate. The parameterization is

$$\begin{aligned}x &= \cos(h) \cos(\theta) \\y &= h \\z &= \cos(h) \sin(\theta),\end{aligned}$$

where the parameter domain  $R$  in the  $h\theta$ -plane consists of  $\frac{\pi}{2} \leq h \leq \frac{\pi}{2}$ ,  $0 \leq \theta \leq 2\pi$ .

1. Calculate  $\vec{r}_h$  and  $\vec{r}_\theta$ .

2. For the line  $0 \leq h \leq \frac{\pi}{2}$ ,  $\theta = \frac{\pi}{4}$  in  $R$ , draw the corresponding points on  $S$ . Do the same for the line  $h = \frac{\pi}{4}$ ,  $0 \leq \theta \leq \frac{\pi}{2}$ . At the intersection point of these two curves on  $S$ , draw the vectors  $\vec{r}_\theta$  and  $\vec{r}_h$ .



(This problem is continued on the next page)

---

(This problem is continued from the previous page.)

3. Given that  $\vec{r}_h \times \vec{r}_\theta = \cos(h) \cos(\theta) \vec{i} + \sin(h) \cos(h) \vec{j} + \cos(h) \sin(\theta) \vec{k}$ , carry out the following:

(a) Check that  $\|\vec{r}_h \times \vec{r}_\theta\| = \cos(h) \sqrt{1 + \sin^2(h)}$ .

(b) Calculate  $\iint_S \sqrt{1 + \sin^2(y)} dS$ .

(c) Calculate  $\iint_S \sin(y) \cos(y) \vec{j} \cdot d\vec{S}$ , where the outward normal is used on  $S$ .

(d) Given that the volume of the football is  $\frac{\pi^2}{2}$ , calculate the flux of the vector field  $(x+y)\vec{i} + (y+z)\vec{j} + (z+x)\vec{k}$  across  $S$ , where the outward normal is used on  $S$ .

**X.** Let  $D$  be a domain in the plane. Define what it means to say that  $D$  is *simply-connected*.  
(3)

**XI.** Let  $T$  be the part of the plane  $x + y + z = 1$  that lies in the first octant, with the upward normal, and let  
(10)  $C$  be its boundary, with the positive orientation.

1. Use Stokes' Theorem to write  $\int_C (xz \vec{i} + 3xy \vec{j} + 3xy \vec{k}) \cdot d\vec{r}$  as a surface integral on  $T$ .

2. Regarding  $T$  as the graph of the function  $z = 1 - x - y$  over a domain  $D$  in the  $xy$ -plane, use the formula  $\iint_S (P \vec{i} + Q \vec{j} + R \vec{k}) \cdot d\vec{S} = \iint_D -P g_x - Q g_y + R dD$  to calculate the surface integral.