

Instructions: Give brief answers, but clearly indicate your reasoning.

$$x = \rho \cos(\theta) \sin(\phi), y = \rho \sin(\theta) \sin(\phi), z = \rho \cos(\phi), dV = \rho^2 \sin(\phi) d\rho d\phi d\theta, \vec{r}_\phi \times \vec{r}_\theta = a \sin(\phi)(x\vec{i} + y\vec{j} + z\vec{k}),$$
$$\|\vec{r}_\phi \times \vec{r}_\theta\| = a^2 \sin(\phi)$$

$$dS = \sqrt{1 + g_x^2 + g_y^2} dD$$

$$dS = \|\vec{r}_u \times \vec{r}_v\| dD$$

$$\iint_S \vec{F} \cdot d\vec{S} = \iint_S \vec{F} \cdot \vec{n} dS$$

$$\iint_S (P\vec{i} + Q\vec{j} + R\vec{k}) \cdot d\vec{S} = \iint_D -P g_x - Q g_y + R dD$$

$$\iint_S \vec{F} \cdot d\vec{S} = \iint_D \vec{F} \cdot (\vec{r}_u \times \vec{r}_v) dD$$

I. A path C is parameterized as a vector-valued function by $\vec{r}(t) = t^2\vec{i} + t\vec{j}$, $1 \leq t \leq 2$. Using this parameterization, evaluate the following line integrals.

1. $\int_C (x/y) dx$

We have $dx = 2t dt$, so $\int_C (x/y) dx = \int_1^2 (t^2/t) 2t dt = \int_1^2 2t^2 dt = 14/3$.

2. $\int_C (x/y) ds$

We have $ds^2 = dx^2 + dy^2 = (2t dt)^2 + (dt)^2 = (4t^2 + 1) dt^2$, so $ds = \sqrt{4t^2 + 1} dt$. So $\int_C (x/y) ds = \int_1^2 (t^2/t) \sqrt{4t^2 + 1} dt = \int_1^2 t \sqrt{4t^2 + 1} dt = (1/8)(2/3)(4t^2 + 1)^{3/2} \Big|_1^2 = (17\sqrt{17} - 5\sqrt{5})/12$.

II. Let $\vec{F}(x, y, z) = 2xy\vec{i} + (x^2 + 2yz)\vec{j} + (y^2 + z)\vec{k}$.

(6)

1. Find a function f such that $\vec{F} = \nabla f$.

We need $f_x = 2xy$, so $f(x, y, z) = x^2y + g(y, z)$ for some function g . We also need $x^2 + 2yz = f_y = x^2 + g_y$, so $g_y = 2yz$ and therefore $g(y, z) = y^2z + h(z)$ and $f(x, y, z) = x^2y + y^2z + h(z)$. Finally, we need $y^2 + z = f_z = y^2 + h'(z)$, so $h'(z) = z$ and therefore $h(z) = z^2/2 + C$. So any f of the form $x^2y + y^2z + z^2/2 + C$ has $\nabla f = \vec{F}$.

2. Calculate $\int_C \vec{F} \cdot d\vec{r}$, where C is given by the parameterization $x = \cos^3(t)$, $y = \cos^4(t)$, $z = \sqrt{\cos(t)}$, $0 \leq t \leq \pi/2$.

We apply the Fundamental Theorem for Line Integrals. The initial point of C is $(1, 1, 1)$, and its terminal point is $(0, 0, 0)$. So for the $f(x, y, z)$ in part 1 (taking $C = 0$) which had $\nabla f = \vec{F}$, we have $\int_C \vec{F} \cdot d\vec{r} = f(0, 0, 0) - f(1, 1, 1) = 0 - 5/2 = -5/2$.

- III.** Let $\vec{F}(x, y)$ be the vector field $\frac{-y}{x^2 + y^2}\vec{i} + \frac{x}{x^2 + y^2}\vec{j}$. Verify by calculation that $\int_C \vec{F} \cdot d\vec{r}$ is not path-
(5) independent on the domain $\{(x, y) \mid (x, y) \neq (0, 0)\}$. (Hint: Consider the line integral of \vec{F} on the unit circle C).

On the unit circle, the unit tangent vector is $\vec{T} = -y\vec{i} + x\vec{j}$, and $x^2 + y^2 = 1$, so we have

$$\int_C \left(\frac{-y}{x^2 + y^2}\vec{i} + \frac{x}{x^2 + y^2}\vec{j} \right) \cdot d\vec{r} = \int_C (-y\vec{i} + x\vec{j}) \cdot \vec{T} ds = \int_C y^2 + x^2 ds = \int_C 1 ds = 2\pi ,$$

since $\int_C 1 ds$ is just the length of C . But when an integral is path-independent, the integral around any closed loop must be 0 (if the integral were path independent, then $\int_C \vec{F} \cdot d\vec{r}$ this would be the same as the integral around the reverse path $-C$, which is -2π).

- IV.** Verify that if $P(x, y, z)\vec{i} + Q(x, y, z)\vec{j} + R(x, y, z)\vec{k}$ is conservative, then $\frac{\partial Q}{\partial z} = \frac{\partial R}{\partial y}$. (Hint: if it is conser-
(4) vative, then it can be written in the form $f_x\vec{i} + f_y\vec{j} + f_z\vec{k}$.)

A conservative vector field can be written in the form $f_x\vec{i} + f_y\vec{j} + f_z\vec{k}$, that is, $Q(x, y, z) = f_y$ and $R(x, y, z) = f_z$. So $\frac{\partial Q}{\partial z} = f_{yz}$ and $\frac{\partial R}{\partial y} = f_{zy}$. By Clairaut's Theorem, these must be equal.

- V.** Suppose that C is a closed loop with no self intersections, bounding a region D .
(5)

1. Explain how one determines the "positive" or "standard" orientation on C .

If you travel along C in the positive direction, you see the region in the plane bounded by C on your left, rather than on your right.

2. State Green's Theorem.

A closed loop C bounds a region R in the plane, and C is given the positive orientation. Green's Theorem says that for functions $P(x, y)$ and $Q(x, y)$,

$$\int_C P(x, y) dx + Q(x, y) dy = \iint_R \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dR .$$

(Alternatively, one can state this in terms of the line integral of a vector field:

$$\int_C (P(x, y)\vec{i} + Q(x, y)\vec{j}) \cdot d\vec{r} = \iint_R \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} dR .)$$

- VI.** Calculate the curl and the divergence of the vector field $\vec{F}(x, y, z) = 3z^2\vec{i} - x\cos(y)\vec{j} + 2xz\vec{k}$.
(5)

$$\text{curl}(3z^2\vec{i} + x\cos(y)\vec{j} + 2xz\vec{k}) = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ \frac{\partial}{\partial x} & \frac{\partial}{\partial y} & \frac{\partial}{\partial z} \\ 3z^2 & -x\cos(y) & 2xz \end{vmatrix} = (0 - 0)\vec{i} - (2z - 6z)\vec{j} + (-\cos(y) - 0)\vec{k} =$$

$$4z\vec{j} - \cos(y)\vec{k} \text{ and } \text{div}(3z^2\vec{i} + x\cos(y)\vec{j} + 2xz\vec{k}) = 0 - x(-\sin(y)) + 2x = 2x + x\sin(y).$$

- VII.** Let S be the portion of the cylinder $x^2 + z^2 = 1$ that lies between the vertical planes $y = 0$ and $y = 2 - x$.
 (5) The surface S is parameterized by $x = \cos(\theta)$, $y = h$, $z = \sin(\theta)$ for $0 \leq \theta \leq 2\pi$ and $0 \leq h \leq 2 - \cos(\theta)$.

1. Calculate \vec{r}_θ and \vec{r}_h .

$$\vec{r}_\theta = -\sin(\theta)\vec{i} + \cos(\theta)\vec{k} \text{ and } \vec{r}_h = \vec{j}.$$

2. Calculate $\vec{r}_h \times \vec{r}_\theta$ and $\|\vec{r}_h \times \vec{r}_\theta\|$.

$$\vec{r}_h \times \vec{r}_\theta = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 0 & 1 & 0 \\ -\sin(\theta) & 0 & \cos(\theta) \end{vmatrix} = \cos(\theta)\vec{i} + \sin(\theta)\vec{k}, \text{ so } \|\vec{r}_h \times \vec{r}_\theta\| = \sqrt{\cos^2(\theta) + \sin^2(\theta)} = \sqrt{1} = 1.$$

- VIII.** Let S be the part of the cone $z = \sqrt{x^2 + y^2}$ that lies between the planes $z = 1$ and $z = 2$. Calculate dS
 (6) in terms of dD , where D is the domain in the xy -plane lying beneath S , and use it to calculate $\iint_S z^2 dS$.

$$\begin{aligned} \text{We calculate } dS &= \sqrt{1 + z_x^2 + z_y^2} dD = \sqrt{1 + \left(\frac{x}{\sqrt{x^2 + y^2}}\right)^2 + \left(\frac{y}{\sqrt{x^2 + y^2}}\right)^2} dD \\ &= \sqrt{1 + \frac{x^2}{x^2 + y^2} + \frac{y^2}{x^2 + y^2}} dD = \sqrt{2} dD, \text{ so } \iint_S z^2 dS = \iint_R (\sqrt{x^2 + y^2})^2 \sqrt{2} dD \\ &= \iint_R r^2 \sqrt{2} r dr d\theta = \int_0^{2\pi} d\theta \int_1^2 \sqrt{2} r^3 dr = 2\pi (16 - 1) \sqrt{2}/4 = 15\pi/\sqrt{2}. \end{aligned}$$

- IX.** Calculate $\iint_S (xy\vec{i} + 4x^2\vec{j} + yz\vec{k}) \cdot d\vec{S}$, where S is the surface $z = xe^y$, $0 \leq x \leq 1$, $0 \leq y \leq 2$.
 (6)

$$\begin{aligned} \text{Using the formula } \iint_S (P\vec{i} + Q\vec{j} + R\vec{k}) \cdot d\vec{S} &= \iint_D -P g_x - Q g_y + R dD, \text{ we have} \\ \iint_S (xy\vec{i} + 4x^2\vec{j} + yz\vec{k}) \cdot d\vec{S} &= \int_D -xy e^y - 4x^2 x e^y + yz dD = \int_D -xy e^y - 4x^2 x e^y + yx e^y dD \\ &= \int_D -4x^3 e^y dD = -\int_0^2 e^y dy \int_0^1 4x^3 dx = 1 - e^2. \end{aligned}$$

- X.** Use Green's Theorem to calculate $\int_C (y^3\vec{i} - x^3\vec{j}) \cdot d\vec{r}$, where C is the circle $x^2 + y^2 = 4$ with the clockwise orientation.
 (6)

$$\begin{aligned} \text{Letting } D \text{ be the unit disk, and noting that } C \text{ has the reverse of the positive orientation, Green's Theo-} \\ \text{rem gives us } \int_C (y^3\vec{i} - x^3\vec{j}) \cdot d\vec{r} &= -\iint_D \frac{\partial(-x^3)}{\partial x} - \frac{\partial(y^3)}{\partial y} dD = \iint_D 3x^2 + 3y^2 dD = \int_0^{2\pi} d\theta \int_0^1 3r^3 dr = \\ &= 2\pi (3/4) \cdot 15 = 24\pi. \end{aligned}$$