

I. Use integration to find a function $f(x, y, z)$ for which $\nabla f = (y + z)\vec{i} + (x + z)\vec{j} + (x + y)\vec{k}$.
(4)

II. Let C be the portion of the circle of radius 2 with center at the origin that lies in the first quadrant $x \geq 0$, $y \geq 0$. By direct calculation using a parameterization of C , evaluate the following line integrals.

1. $\int_C x^2 y \, ds$

2. $\int_C xy \, dy$

3. $\int_C (x\vec{i} + y\vec{j}) \cdot d\vec{r}$


III. Use the Fundamental Theorem of Calculus to carry out a partial calculation of $\iint_R \frac{\partial Q}{\partial y} \, dA$, where R is the rectangle $1 \leq x \leq 3$, $2 \leq y \leq 4$, and $Q(x, y)$ is a function of x and y .
(3)

- IV. Use Green's Theorem to calculate $\int_C 3xy dx + 5x^2y^2 dy$, where C is the triangle with vertices $(0, 0)$, $(1, 0)$, and $(1, 1)$.

- V. Calculate the curl and the divergence of the vector field $x^2\vec{i} + y^2\vec{j} - xyz\vec{k}$.

- VI. The figure to the right shows a vector field $\vec{F} = P\vec{i} + Q\vec{j}$ and three oriented arcs.

- Near each arc, write a small "+" if the line integral of \vec{F} along that arc appears to be positive, a "-" if it appears to be negative, and a "0" if it appears to be 0.
- Does it appear that $\frac{\partial P}{\partial x}$ is positive, negative, or 0?
- Does it appear that $\frac{\partial Q}{\partial y}$ is positive, negative, or 0?
- Does it appear that $\text{div}(\vec{F})$ is positive, negative, or 0?



VII. Let S be the surface given by $x = u \cos(v)$, $y = u \sin(v)$, and $z = u$, where the domain of the parameterization is the rectangle $0 \leq u \leq 1$ and $0 \leq v \leq 2\pi$.

1. Calculate \vec{r}_u , \vec{r}_v , $\vec{r}_u \times \vec{r}_v$, and $\|\vec{r}_u \times \vec{r}_v\|$.


2. Sketch the domain R in the uv -plane. Tell the points in R where locally the parameterization *neither stretches nor contracts area*.

3. Find an equation in x , y , and z satisfied by all points in the surface (hint: start by calculating $x^2 + y^2$).

VIII. Let $f(x, y, z) = \sin(x^2 + y^2 + z)$. Let C_1 be the line segment from $(0, 0, 0)$ to $(1, 1, 0)$, and let C_2 be the curve on the surface $z = e^{xy}$ that lies directly above C_1 . Calculate $\int_{C_1} \nabla f \cdot d\vec{r}$ and $\int_{C_2} \nabla f \cdot d\vec{r}$.

- IX.** Let C be the unit circle in the xy -plane and let \vec{T} be its unit tangent vector. Suppose that a certain vector field \vec{F} has the property that each point (x, y) in C , $\vec{F} \cdot \vec{T} = \pi$. Find $\int_C \vec{F} \cdot d\vec{r}$.

- X.** The figure below shows four regions in the plane. Below each region, write a very small letter m if the region is simply connected, and a very small letter n if the region is not simply-connected. The three dots on the last region means that the region continues to the right forever.



- XI.** Find a vector field \vec{F} in the plane so that if C is any path which does not pass through the origin, and C starts at P and ends at Q , then $\int_C \vec{F} \cdot d\vec{r}$ equals the distance from Q to the origin, minus the distance from P to the origin.

- XII.** Give an example of a 2-dimensional vector field $P\vec{i} + Q\vec{j}$ which is not conservative but which does satisfy the condition $\frac{\partial Q}{\partial x} = \frac{\partial P}{\partial y}$. You do not need to verify these properties, just write down the vector field.