


Problem 1. The purpose of this problem is to construct and study the Newton's divided difference form of the interpolating polynomial,

$$P_n(x) = f[x_0] + \sum_{k=1}^n f[x_0, \dots, x_k] \prod_{j=0}^{k-1} (x - x_j),$$

to the function $f(x) = \cos(\pi x)$. The points $x_i, i = 0, 1, 2, 3$ used to construct the interpolating polynomial are given in the table below.

The graphs of the function $f(x)$ and the interpolating polynomials $P_0(x), P_1(x), P_2(x)$, and $P_3(x)$ are plotted in the figure below (the graph of $f(x)$ is plotted with a dashed line).



- (a) Compute the missing entries in the divided differences table below. Write your calculations clearly and leave the coefficients in symbolic form (i.e., do not compute the numerical values of things like $12(8\sqrt{2} - 11)$).

x_i	0 th order	1 st order	2 nd order	3 rd order
$x_0 = 0$	$f[x_0] = 1$			
		$f[x_0, x_1] = ?$		
$x_1 = \frac{1}{3}$	$f[x_1] = \frac{1}{2}$		$f[x_0, x_1, x_2] = ?$	
		$f[x_1, x_2] = ?$		$f[x_0, x_1, x_2, x_3] = 12(8\sqrt{2} - 11)$
$x_2 = \frac{1}{2}$	$f[x_2] = 0$		$f[x_1, x_2, x_3] = 12(2\sqrt{2} - 3)$	
		$f[x_2, x_3] = -2\sqrt{2}$		
$x_3 = \frac{1}{4}$	$f[x_3] = ?$			

- (b) Write down the interpolating polynomial $P_0(x)$ based on the values in the divided differences table above.
- (c) Similarly to part (b), write down the interpolating polynomial $P_1(x)$ based on the values in the divided differences table above.
- (d) Similarly to part (b), write down the interpolating polynomial $P_2(x)$ based on the values in the divided differences table above. Do *not* expand it – just substitute the coefficients in the Newton’s divided difference interpolating polynomial with the corresponding entries from the table.
- (e) Similarly to part (b), write down the interpolating polynomial $P_3(x)$ based on the values in the divided differences table above. Do *not* expand the polynomial!

Problem 2. Use Newton’s forward-difference formula to construct interpolating polynomials for the function $f(x) = \cos(\pi x)$ based on values of $f(x)$ at equally spaced points (note that the points x_0, x_1, x_2 and x_3 below are equally spaced with $h = \frac{1}{6}$). First write the polynomial in the form

$$P_n(x) = f(x_0) + \sum_{k=1}^n \binom{s}{k} \Delta^k f(x_0),$$

where $x = x_0 + sh$, i.e., $s = \frac{x-x_0}{h}$.

Fill the missing entries in the forward differences table below, and find the following:

- (a) the degree-1 forward difference polynomial $P_1(x)$ based on the points $(x_0, f(x_0)) = (0, 1)$ and $(x_1, f(x_1)) = (\frac{1}{6}, \frac{\sqrt{3}}{2})$;
- (b) the degree-2 forward difference polynomial $P_2(x)$ based on the points $(x_0, f(x_0))$, $(x_1, f(x_1))$, and $(x_2, f(x_2)) = (\frac{1}{3}, ?)$;
- (c) the degree-3 forward difference polynomial $P_3(x)$ based on the points $(x_0, f(x_0))$, $(x_1, f(x_1))$, $(x_2, f(x_2))$, and $(x_3, f(x_3)) = (\frac{1}{2}, ?)$.
- (d) In Mathematica (or any other computer system), plot the graphs of $\cos(\pi x)$, $P_1(x)$, $P_2(x)$, and $P_3(x)$.

Use the forward differences table below, some of whose entries are given. Do *not* compute the numerical values (i.e., leave the square roots, as in the table below, and, more importantly, leave the polynomial in Newton’s forward-difference form (i.e., do *not* expand it).

x_i	0 th order	1 st order	2 nd order	3 rd order
$x_0 = 0$	$f(x_0) = 1$	$\Delta f(x_0) = ?$		
$x_1 = \frac{1}{6}$	$f(x_1) = \frac{\sqrt{3}}{2}$	$\Delta f(x_1) = ?$	$\Delta^2 f(x_0) = ?$	
$x_2 = \frac{1}{3}$	$f(x_2) = ?$	$\Delta f(x_2) = -\frac{1}{2}$	$\Delta^2 f(x_1) = \frac{\sqrt{3}-2}{2}$	$\Delta^3 f(x_0) = \frac{3\sqrt{3}-5}{2}$
$x_3 = \frac{1}{2}$	$f(x_3) = ?$			

Problem 3. Consider the function $f(x) = \sqrt{x}$. We want to find a cubic polynomial $S(x)$ that interpolates the function $f(x)$ on the interval $x \in [1, 4]$ with clamped boundary conditions at both ends.

- Clearly, $S(x)$ must have the same values as $f(x)$ at the points $x_0 = 1$ and $x_1 = 4$. What are the clamped boundary conditions for $S'(1)$ and $S'(4)$?
- Write the interpolating polynomial $S(x)$ in the form

$$S(x) = a + b(x - 1) + c(x - 1)^2 + d(x - 1)^3 ,$$

and impose the conditions formulated in part (a) to find the coefficients of $S(x)$. You should obtain that $c = -\frac{1}{12}$ and $d = \frac{1}{108}$.

- Apply Theorem 3.13 on page 152 to give a rigorous upper bound on the interpolation error. Compare this numerical value with the actual value of the error, which is about 0.013.
- The function $S(x)$ that you found in (a) is good not only to approximate the values of the function $f(x)$, but also to approximate the values of the integrals and some derivatives of $f(x)$. Find the numerical value of $\int_1^3 S(x) dx$ and compare it with the exact value, $\int_1^3 f(x) dx$; find the absolute and the relative errors.

Problem 4. The polynomials

$$S_0(x) = 4 + 4(x - 1) + 13(x - 1)^2 - 9(x - 1)^3 , \quad x \in [1, 2] ,$$

$$S_1(x) = a + b(x - 2) + c(x - 2)^2 + d(x - 2)^3 , \quad x \in [2, 3]$$

form the *clamped* cubic spline interpolant for some function $f(x)$ that is known to satisfy $f'(1) = -f'(3)$.

- Use this information to find the values of the coefficients a , b , c , and d .
- Compute $S'(2.5)$.